## Author index

Adamson, P., see Romero, I.A. (105) 47 Ajpru, S., McArthur, A.J., Piggins, H.D. and Sugden, D.

Identification of PAC1 receptor isoform mRNAs by real-time PCR in rat suprachiasmatic nucleus (105) 29

Amos, C.L., see Romero, I.A. (105) 47
Andersen, T.E., Finsen, B., Goffinet, A.M.,
Issinger, O.-G. and Boldyreff, B.
A reeler mutant mouse with a new,
spontaneous mutation in the reelin
gene (105) 153

Andria, M.L. and Simon, E.J.
Erratum to: Identification of a
neurorestrictive suppressor element
(NRSE) in the human μ-opioid
receptor gene. [Molecular Brain
Research 91 (2001) 73–80] (105)
161

Antonio González-Barrios, J., see Navarro-Quiroga, I. (105) 86

Bales, K.R., see Gao, F. (105) 108 Barron-Moreno, F., see Navarro-Quiroga, I. (105) 86

Black, J.A., Renganathan, M. and Waxman, S.G.
 Sodium channel Na<sub>v</sub>1.6 is expressed along nonmyelinated axons and it contributes to conduction (105) 19

Boldyreff, B., see Andersen, T.E. (105) 153 Bridges, R.S., see Mann, P.E. (105) 136

Chen, X., see Gao, F. (105) 108 Cho, H.J., see Kim, D.S. (105) 146 Choi, J.O., see Kim, D.S. (105) 146

Daudet, N., Ripoll, C., Molès, J.-P. and Rebillard, G. Expression of members of *Wnt* and *Frizzled* gene families in the postnatal rat cochlea (105) 98

Dodel, R.C., see Gao, F. (105) 108 Du, Y., see Gao, F. (105) 108

Fahnestock, M., see Yu, G. (105) 115 Fahrenkrug, J., see Nielsen, H.S. (105) 79 Farlow, M.R., see Gao, F. (105) 108 Finsen, B., see Andersen, T.E. (105) 153

Gao, F., Bales, K.R., Dodel, R.C., Liu, J., Chen, X., Hample, H., Farlow, M.R., Paul, S.M. and Du, Y. NF-κB mediates IL-1β-induced synthesis/release of α<sub>2</sub>-macroglobulin in a human glial cell line (105) 108 Georg, B., see Nielsen, H.S. (105) 79 Gietzen, D.W., see Sharp, J.W. (105) 11 Goffinet, A.M., see Andersen, T.E. (105) 153 González-Bernal, V., see Navarro-Quiroga, I.

(105) 86 Greenwood, J., see Romero, I.A. (105) 47

Hample, H., see Gao, F. (105) 108
Hannibal, J., see Nielsen, H.S. (105) 79
Hoshikawa, M., Yonamine, A., Konishi, M. and Itoh, N.
FGF-18 is a neuron-derived glial cell growth factor expressed in the rat brain during early postnatal

Issinger, O.-G., see Andersen, T.E. (105) 153 Itoh, N., see Hoshikawa, M. (105) 60

development (105) 60

Kaplan, I.V., see Mower, G.D. (105) 157 Kim, D.S., Choi, J.O., Rim, H.D. and Cho, H.J.

Downregulation of voltage-gated potassium channel  $\alpha$  gene expression in dorsal root ganglia following chronic constriction injury of the rat sciatic nerve (105) 146

Kojima, N., see Stork, O. (105) 126 Konishi, M., see Hoshikawa, M. (105) 60 Kume, N., see Stork, O. (105) 126

Leonova, E.V. and Lomax, M.I.

Expression of the mouse *Macf2* gene during inner ear development (105)

Liu, J., see Gao, F. (105) 108 Lomax, M.I., see Leonova, E.V. (105) 67

Magrum, L.J., see Sharp, J.W. (105) 11 Mann, P.E. and Bridges, R.S.

Prolactin receptor gene expression in the forebrain of pregnant and lactating rats (105) 136

Martinez-Arguelles, D.B., see Navarro-Quiroga, I. (105) 86

Martinez-Fong, D., see Navarro-Quiroga, I. (105) 86

McArthur, A.J., see Ajpru, S. (105) 29 Molès, J.-P., see Daudet, N. (105) 98 Mower, G.D. and Kaplan, I.V. Immediate early gene expression in the visual cortex of normal and dark reared cats: differences between *fos* and *egr-1* (105) 157

Navarro-Quiroga, I., Antonio González-Barrios, J., Barron-Moreno, F., González-Bernal, V., Martinez-Arguelles, D.B. and Martinez-Fong, D. Improved neurotensin-vector-mediated gene transfer by the coupling of hemagglutinin HA2 fusogenic peptide and Vp1 SV40 nuclear localization signal (105) 86

Nielsen, H.S., Georg, B., Hannibal, J. and Fahrenkrug, J.

Homer-1 mRNA in the rat suprachiasmatic nucleus is regulated differentially by the retinohypothalamic tract transmitters pituitary adenylate cyclase activating polypeptide and glutamate at time points where light phase-shifts the endogenous rhythm (105) 79

Obata, K., see Stork, O. (105) 126 Oberdick, J., see Zhang, X. (105) 1

Paul, S.M., see Gao, F. (105) 108 Piggins, H.D., see Ajpru, S. (105) 29

Rebillard, G., see Daudet, N. (105) 98
Renganathan, M., see Black, J.A. (105) 19
Rim, H.D., see Kim, D.S. (105) 146
Ripoll, C., see Daudet, N. (105) 98
Romero, I.A., Amos, C.L., Greenwood, J. and Adamson, P.

Ezrin and moesin co-localise with ICAM-1 in brain endothelial cells but are not directly associated (105) 47

Sharp, J.W., Magrum, L.J. and Gietzen, D.W. Role of MAP kinase in signaling indispensable amino acid deficiency in the brain (105) 11

Simon, E.J., see Andria, M.L. (105) 161 Standifer, K.M., see Thakker, D.R. (105) 38 Stork, O., Kojima, N., Stork, S., Kume, N. and Obata, K.

> Resistance to alcohol withdrawalinduced behaviour in Fyn transgenic mice and its reversal by ifenprodil (105) 126

Stork, S., see Stork, O. (105) 126 Sugden, D., see Ajpru, S. (105) 29

Thakker, D.R. and Standifer, K.M.
Orphanin FQ/nociceptin blocks
chronic morphine-induced tyrosine
hydroxylase upregulation (105) 38

Waxman, S.G., see Black, J.A. (105) 19

Yonamine, A., see Hoshikawa, M. (105) 60 Yu, G. and Fahnestock, M. Differential expression of nerve growth factor transcripts in glia and neurons and their regulation by transforming growth factor-β1 (105)

Zhang, H., see Zhang, X. (105) 1

Zhang, X., Zhang, H. and Oberdick, J.
Conservation of the developmentally regulated dendritic localization of a Purkinje cell-specific mRNA that encodes a G-protein modulator: comparison of rodent and human Pcp2(L7) gene structure and expression (105) 1

.8 0.